## **REVIEW**

## Qualitative and Quantitative Intravaginal Targeting: Key to Anti-HIV-1 Microbicide Delivery from Test Tube to In Vivo Success

Published online in Wiley Online Library (wileyonlinelibrary.com). DOI 10.1002/jps.23098

## VINESS PILLAY,<sup>1</sup> FELIX MASHINGAIDZE,<sup>1</sup> YAHYA E. CHOONARA,<sup>1</sup> LISA C. DU TOIT,<sup>1</sup> ECKHART BUCHMANN,<sup>2</sup> VINESH MAHARAJ,<sup>3</sup> VALENCE M. K. NDESENDO,<sup>4</sup> PRADEEP KUMAR<sup>1</sup>

<sup>1</sup>Department of Pharmacy and Pharmacology, Faculty of Health Sciences, University of the Witwatersrand, Parktown, 2193, Johannesburg, South Africa

<sup>2</sup>Department of Obstetrics and Gynecology, Faculty of Health Sciences, University of the Witwatersrand, Parktown, 2193, Johannesburg, South Africa

<sup>3</sup>Department of Biosciences, Council for Scientific and Industrial Research, Brummeria, Pretoria, South Africa <sup>4</sup>School of Pharmacy and Pharmaceutical Sciences, St. John's University of Tanzania, Dodoma, Tanzania

**ABSTRACT:** The past decade has seen several effective anti-HIV-1 agent discoveries, yet microbicides continue to disappoint clinically. Our review expounds the view that unsatisfactory microbicide failures may be a result of inefficient delivery systems employed. We hereby propose a thorough scientific qualitative and quantitative investigation of important aspects involved in HIV-1 transmission as a prerequisite for microbicide delivery. Intravaginal targeting of HIV-1 increases the chances of microbicide success, wherein vaginal microenvironmental factors including pH should be maintained at HIV-1 prohibitive acidic levels simultaneously to ward off other sexually transmitted diseases, which compromise vaginal epithelial barrier properties. Furthermore, choice of receptors to target both on HIV-1 and on target cells is vital in deterring transmission. Appropriate modeling of virus–target cell interactions as well as targeting early stages of the HIV-1 infection accompanied by computation and delivery of appropriate microbicide quantities could revolutionize microbicide research, ultimately delivering a female-controlled HIV-1 prevention modality appropriately.